



COMPREHENSIVE TOBACCO PREVENTION AND CESSATION PROGRAMS EFFECTIVELY REDUCE TOBACCO USE

Tobacco control programs play a crucial role in the prevention of many chronic conditions such as cancer, heart disease, and respiratory illness. Comprehensive tobacco prevention and cessation programs prevent kids from starting to smoking, help adult smokers quit, educate the public, the media and policymakers about policies that reduce tobacco use, address disparities, and serve as a counter to the ever-present tobacco industry.

Recommendations for state tobacco prevention and cessation programs are best summarized in the Center for Disease Control and Prevention's *Best Practices for Comprehensive Tobacco Control Programs*. In this guidance document, CDC recommends that states establish tobacco control programs that are comprehensive, sustainable, and accountable and include state and community interventions, public education interventions, cessation programs, surveillance and evaluation and administration and management.¹

The empirical evidence regarding the effectiveness of comprehensive tobacco prevention and cessation programs is vast and growing. There is more evidence than ever before that tobacco prevention and cessation programs work to reduce smoking, save lives and save money. In 2007, the Institute of Medicine and the President's Cancer Panel all issued landmark reports that concluded there is overwhelming evidence that state comprehensive state tobacco control programs substantially reduce tobacco use and recommended that every state fund such programs at CDC-recommended levels.²

Data from numerous states that have implemented programs consistent with CDC guidelines show significant reductions in youth and adult smoking. The most powerful evidence, however, comes from national studies that look across states and control for as many of the relevant confounding factors as possible. These rigorous studies consistently show effects of tobacco prevention and cessation programs.

A recent study published in the *American Journal of Public Health*, examined state tobacco prevention and cessation funding levels from 1995 to 2003 and found that the more states spent on these programs, the larger the declines they achieved in adult smoking, even when controlling for other factors such as increased tobacco prices. The researchers also calculated that if every state had funded their programs at the levels recommended by the U.S. Centers for Disease Control (CDC) during that period, there would have been between 2.2 million and 7.1 million fewer smokers in the United States by 2003.³ The Campaign for Tobacco-Free Kids estimates that such smoking declines would have saved between 700,000 and 2.2 million lives as well as between \$20 billion and \$67 billion in health care costs.

The study described above adds to earlier research, using similar methods, which demonstrated the same type of relationship between program spending and youth smoking declines. A 2005 study concluded that if every state had spent the minimum amount recommended by the CDC for tobacco prevention, youth smoking rates nationally would have been between three and 14 percent lower during the study period, from 1991 to 2000. Further, if every state funded tobacco prevention at CDC minimum levels, states would prevent nearly two million kids alive today from becoming smokers, save more than 600,000 of them from premature, smoking-caused deaths, and save \$23.4 billion in long-term, smoking-related health care costs.⁴

A 2003 study published in the *Journal of Health Economics* found that states with the best funded and most sustained tobacco prevention programs during the 1990s – Arizona, California, Massachusetts and Oregon – reduced cigarette sales more than twice as much as the country as a whole (43 percent compared to 20 percent). This study, the first to compare cigarette sales data from all the states and to isolate the impact of tobacco control program expenditures from other factors that affect cigarette sales, demonstrates that the more states spend on tobacco prevention, the greater the reductions in smoking, and the longer states invest in such programs, the larger the impact. The study concludes that cigarette sales would have declined by 18 percent instead of nine percent between 1994 and 2000 had all states fully funded tobacco prevention programs.⁵

A 2006 study published in the *American Journal of Health Promotion* provides further evidence of the effectiveness of comprehensive tobacco control programs and tobacco control policies. The study's findings suggest that well-funded tobacco control programs combined with strong tobacco control policies increase cessation rates. Quit rates in communities that experienced both policy and programmatic interventions were higher than quit rates in communities that had only experienced policy interventions (excise tax increases or secondhand smoke regulations). This finding supports the claim that state-based tobacco control programs can accelerate adult cessation rates in the population and have an effect beyond that predicted by tobacco-control policies alone.⁶

Data from numerous states provide additional evidence of the effectiveness of comprehensive tobacco prevention and cessation programs. States that have implemented comprehensive programs have achieved significant reductions in tobacco use among both adults and youth. The experiences in states from around the country who have invested in comprehensive prevention programs establish the following key points:

- When adequately funded, comprehensive state tobacco prevention programs quickly and substantially reduce tobacco use, save lives, and cut smoking-caused costs.
- State tobacco prevention programs must be insulated against the inevitable attempts by the tobacco industry to reduce program funding and otherwise interfere with the programs' successful operation.
- The programs' funding must be sustained over time both to protect initial tobacco use reductions and to achieve further cuts.
- When program funding is cut, progress in reducing tobacco use erodes, and the state suffers from higher levels of smoking and more smoking-caused deaths, disease, and costs.

Unfortunately, many states faced with budget difficulties have recently made the penny-wise but pound-foolish decision to slash the funding of even the most effective tobacco control programs, which will cost lives and money.

Program Success – California

In 1988, California voters approved Proposition 99, a ballot initiative that increased state cigarette taxes by 25 cents per pack, with 20 percent of the new revenues (over \$100 million per year) earmarked for health education against tobacco use. California launched its new Tobacco Control Program in Spring 1990. Despite increased levels of tobacco marketing and promotion, a major cigarette price cut in 1993, tobacco company interference with the program, and periodic cuts in funding, the program has still reduced tobacco use and its attendant devastation substantially.

- California's comprehensive approach has reduced adult smoking significantly. Adult smoking declined by 43 percent from 1988 to 2007, from 24.2 percent to 13.8 percent.⁷ If every state had California's current smoking rate, there would be more than 16 million fewer smokers in the United States.
- Since the passage of Proposition 99, between 1988 and 2003, cigarette consumption in California declined by 60 percent, compared to just 38 percent for the country as a whole.⁸ Even after the tobacco industry's successful efforts to reduce the state's tobacco prevention funding, cigarette consumption still declined more in California than in the rest of the country.⁹
- In the 10 years following the passage of Proposition 99, adult smoking in California declined at twice the rate it declined in the previous decade.¹⁰

* This factsheet focuses on the extensive public health benefits obtained by state tobacco prevention programs. Other Campaign factsheets show that these programs also reduce smoking-caused costs, including those incurred by state Medicaid programs. See, e.g., TFK Factsheet, *Comprehensive Statewide Tobacco-Prevention Programs Save Money*, <http://tobaccofreekids.org/research/factsheets/pdf/0168.pdf>.

- Between 1988 and 2003, lung and bronchus cancer rates in California declined at three times the rate of decline as the rest of the U.S.¹¹ Surveillance, Epidemiology, and End Results (SEER) data associated lower lung cancer incidence with California's program.¹²
- In California, from 1996 to 2006, smoking declined by 45 percent among eighth grade students and by 46 percent among tenth grade students. From 2000 to 2006, smoking prevalence decreased by more than 20 percent among twelfth grade students.¹³

The California tobacco control program produced much larger smoking reductions in the early years, when it was funded at its highest levels, than during subsequent years, when the state cut its funding. For example, when California cut the program's funding in the mid 1990s, its progress in reducing adult and youth smoking rates stalled, but it got back on track when program funding was partially restored.¹⁴

Program Success – Washington

The Washington State Tobacco Prevention and Control program was implemented in 1999 after the state Legislature set aside money from the Master Settlement Agreement to create a Tobacco Prevention and Control Account. Tobacco prevention and control received additional funds in 2001 when the state's voters passed a cigarette tax increase that dedicated a portion of the new revenue to tobacco prevention and cessation. According to a recent study in CDC's peer-reviewed journal, *Preventing Chronic Disease*, although Washington made progress in implementing tobacco control policies between 1990 and 2000, smoking prevalence did not decline significantly until after substantial investment was made in the state's comprehensive tobacco control program.¹⁵ As the data below demonstrate, Washington's comprehensive program is working.

- Since the program began, Washington's tobacco prevention efforts have cut smoking by 60 percent among sixth graders, 58 percent among eighth graders, 40 percent among tenth graders, and 43 percent among twelfth graders. Because of these declines, there are 65,000 fewer youth smokers in Washington.¹⁶
- Since the tobacco control program was implemented, adult smoking has declined by 24 percent, from 22.4 percent in 1999 to 16.5 percent in 2007, one of the lowest smoking rates in the country.¹⁷ Washington's dramatic decline in adult smoking translates to more than 240,000 fewer smokers in the state, saving about \$2.1 billion in future health care costs.¹⁸

Program Success – New York

New York began implementing a comprehensive state tobacco control program in 2000 with funds from the Master Settlement Agreement and revenue from the state cigarette tax. As the data below demonstrate, New York's comprehensive approach is working. While declines in youth smoking nationally have slowed, New York's rates continue to decline steadily.

- Between 2000 and 2006, smoking among middle school students declined by 61 percent, (from 10.5 percent to 4.1 percent), and smoking among high school students declined by 40 percent, (from 27.1 percent to 16.3 percent).¹⁹
- Between 2000 and 2006, adult smoking declined by 15 percent, from 21.6 percent to 18.3 percent.²⁰

Program Success – Maine

In 1997, Maine increased its cigarette excise tax and used a portion of those funds to establish a comprehensive tobacco prevention program known as the Partnership for a Tobacco-Free Maine. Maine has subsequently augmented its program with proceeds from the 1998 state tobacco settlement, which also resulted in a further increase in cigarette prices (the state also raised cigarette taxes again in 2001, to \$1.00 per pack, and in 2005 to \$2.00 per pack). Prior to launching this effort, Maine had one of the highest youth smoking rates in the country. Now, it has one of the lowest.

- Smoking among Maine's high school students declined a dramatic 64 percent between 1997 and 2007, falling from 39.2 percent to 14 percent. Smoking among Maine's middle school students declined by 71 percent, from 21 percent to 6 percent, over the same time period.²¹ The Maine Department of Health (DOH) has calculated that, as a result of these declines, there are now more than 26,000 fewer youth smokers in Maine and more than 14,000 youth will be saved from premature, smoking-caused deaths. Based on estimates that smokers, on average, have \$16,000 more in lifetime health care costs than non-smokers, the DOH calculated that these declines will save Maine more than \$416 million in long-term health care costs.

Program Success – Indiana

In 2000, Indiana implemented a comprehensive tobacco prevention and cessation program with revenue received from the state's tobacco settlement. Indiana's program is modeled after other comprehensive programs that have been successful in reducing tobacco use. Indiana's program includes public education efforts, a counter-marketing campaign, community and school-based programs, and enforcement initiatives.²²

- Between 2000 and 2006, smoking among high school students declined by 25 percent, (from 32.0 percent to 23.9 percent).
- Smoking among middle school students declined by 22 percent, from 10 percent to 7.8 percent, over this same time period.

Program Success – An Experiment in Texas

Rather than using settlement money to fund a comprehensive statewide tobacco prevention program, the state of Texas decided to use a small portion of its tobacco settlement money to test tobacco prevention interventions of varying intensity and comprehensiveness in selected parts of the state. Not surprisingly, this experiment found that the largest effects on both youth smoking rates occurred in those areas where comprehensive programs were implemented and sustained. Data show that youth smoking in the comprehensive program area decreased at more than four times the state rate of decline.²³

- Between 2000 and 2005, smoking among high school students dropped by 46 percent, from 34.2 percent to 18.3 percent, in the Beaumont/Port Arthur comprehensive program area. Statewide, youth smoking only declined by 9.3 percent, from 24.7 percent in 2001 to 22.4 percent in 2004.
- From 2000 to 2005, current cigarette use among middle school students decreased by 34 percent (from 17 percent to 11.2 percent) in the Beaumont/Port Arthur comprehensive program area. Statewide, smoking among middle school students actually increased by 2 percent, from 10.2 percent to 10.4 percent, between 2001 and 2004.

Program Success -- Massachusetts

In 1992, Massachusetts voters approved a referendum that increased the state cigarette tax by 25 cents per pack. Part of the new tax revenues was used to fund the Massachusetts Tobacco Control Program (MTCP), which began in 1993. As in California, despite some reductions in funding encouraged by the tobacco industry, the program achieved considerable success until its funding was cut by more than 90 percent in 2003. Data from 2000 demonstrate that the program was successful in reducing tobacco use among both children and adults.

- Massachusetts cigarette consumption declined by 36 percent between 1992 and 2000, compared to a decrease of just 16 percent in the rest of the country (excluding California).²⁴
- From 1995 to 2001, current smoking among Massachusetts high school students dropped by 27 percent (from 35.7 percent to 26 percent), while the nationwide rate dropped by 18 percent (34.8 percent to 28.5 percent)²⁵

- Between 1993 and 2000, adult smoking prevalence dropped from 22.6 percent to 17.9 percent, resulting in 228,000 fewer smokers.²⁶ Nationally, smoking prevalence dropped by just seven percent over this same time period.²⁷
- Between 1990 and 1999, smoking among pregnant women in Massachusetts declined by more than 50 percent (from 25 percent to 11 percent). Massachusetts had the greatest percentage decrease of any state over the time period (the District of Columbia had a greater percent decline).²⁸

Despite the considerable success achieved in Massachusetts, funding for the state's tobacco prevention and cessation program was cut by 95 percent – from a high of approximately \$54 million per year to just \$2.5 million in FY2004, although funding for the program has increased slightly in recent years. These drastic reductions in the state's investments to prevent and reduce tobacco use will translate directly into higher smoking rates, especially among kids, and more smoking-caused disease, death, and costs. In fact, a study released by the Massachusetts Association of Health Boards shows that the Massachusetts program funding cuts have already been followed by an alarming increase in illegal sales of tobacco products to children.²⁹

- Between 2002 and 2003, cigarette sales to minors increased by 74 percent, from eight percent to 13.9 percent in communities that lost a significant portion of their enforcement funding.
- Over the same time period, cigarette sales to minors increased by 98 percent in communities that lost all of their local enforcement funding.
- Between 1992 and 2003, per capita cigarette consumption declined at a higher rate in Massachusetts as it did in the country as a whole (47 percent v. 28 percent). However, from 2003 to 2006, Massachusetts' per capita cigarette consumption declined a mere seven percent (from 47.5 to 44.1 packs per capita), while the U.S. average cigarette consumption declined by ten percent (from 67.9 to 61.1 packs per capita). Most recently, between 2005 and 2006, Massachusetts' per capita cigarette consumption *increased* by 3.2 percent (from 42.7 to 44.1 packs per capita), while nationwide, per capita consumption *declined* by 3.5 percent (from 63.3 to 61.1 packs per capita).³⁰

Campaign for Tobacco-Free Kids, January 26, 2009 / Meg Riordan

¹ Centers for Disease Control and Prevention (CDC), *Best Practices for Comprehensive Tobacco Control Programs*, Atlanta, GA: U.S. Department of Health and Human Services (HHS), October 2007, http://www.cdc.gov/tobacco/tobacco_control_programs/stateandcommunity/best_practices. v

² Institute of Medicine, *Ending the Tobacco Problem: A Blueprint for the Nation*, National Academy of Sciences, 2007; *Promoting Healthy Lifestyles: Policy, Program and Personal Recommendations for Reducing Cancer Risk*, 2006-2007 Annual Report, President's Cancer Panel; See also, Institute of Medicine, *State Programs Can Reduce Tobacco Use*, National Academy of Sciences, 2000; HHS, *Reducing Tobacco Use: A Report of the Surgeon General*, 2000.

³ Farrelly, MC, et al., "The Impact of Tobacco Control Programs on Adult Smoking," *American Journal of Public Health* 98:304-309, February 2008.

⁴ Tauras, JA, et al., "State Tobacco Control Spending and Youth Smoking," *American Journal of Public Health* 95:338-344, February 2005.

⁵ Farrelly, MC, et al., "The Impact of Tobacco Control Program Expenditures on Aggregate Cigarette Sales: 1981-2000," *Journal of Health Economics* 22:843-859, 2003.

⁶ Hyland, A, et al., "State and Community Tobacco-Control Programs and Smoking – Cessation Rates Among Adult Smokers: What Can We Learn From the COMMIT Intervention Cohort?" *American Journal of Health Promotion* 20(4):272, April/March 2006.

⁷ Adult Smoking Prevalence, California Department of Health Services, Tobacco Control Section, 2007 <http://www.dhs.ca.gov/tobacco>. See also, Overview of Evaluation in the California Tobacco Control Program; Warner, Kenneth E, et al., "Tobacco Control Success vs Demographic Destiny: Examining the Causes of the Low Smoking Prevalence in California," *Am J Public Health* 98: 268-269, February 2008.

⁸ Cigarette Consumption, California Department of Health Services, Tobacco Control Section, 2005. <http://www.dhs.ca.gov/tobacco>.

-
- ⁹ Pierce, JP, et al., "Has the California Tobacco Control Program Reduced Smoking?," *Journal of the American Medical Association* 280(10), September 9, 1998.
- ¹⁰ *California's Tobacco Control Program: Preventing Tobacco Related Disease and Death*; Tobacco Control Section, California Department of Health Services, April 3, 1998.
- ¹¹ California Department of Health Services, Tobacco Control Section, California Tobacco Control Update, 2006, <http://www.cdph.ca.gov/programs/tobacco/Documents/CTCPUpdate2006.pdf>; CDC, presentation at the National Conference on Tobacco or Health, 2007, Minneapolis, MN.
- ¹² Cancer Surveillance Section, California Department of Health Services. Unpublished data. See also, California Department of Health Services, Tobacco Control Section, California Tobacco Control Update, 2004, <http://www.dhs.ca.gov/tobacco/documents/pubs/2004TCSupdate.pdf>.
- ¹³ California Department of Health Services, Tobacco Control Section, Youth Smoking, 2008 <http://www.cdph.ca.gov/programs/tobacco/Documents/CTCPFactShYouthSmoking2008.pdf>.
- ¹⁴ Pierce, JP, et al., "Has the California Tobacco Control Program Reduced Smoking?," *Journal of the American Medical Association* 280(10):893-899, September 9, 1998.
- ¹⁵ Dilley JA, et al., "Effective tobacco control in Washington State: A smart investment for healthy futures," *Preventing Chronic Disease* 4(3), July 3, 2007, http://www.cdc.gov/pcd/issues/2007/jul/06_0109.htm.
- ¹⁶ Washington State Department of Health, Tobacco Prevention and Control Program, Progress Report, March 2007. Data are from 2006 Healthy Youth Survey, <http://www.doh.wa.gov/Tobacco/program/reports/tpcp07progrpt.pdf>.
- ¹⁷ CDC, *Behavioral Risk Factor Surveillance System* (BRFSS), <http://apps.nccd.cdc.gov/brfss/display.asp?cat=TU&yr=2006&qkey=4396&state=WA>.
- ¹⁸ Dilley JA, et al., "Effective tobacco control in Washington State: A smart investment for healthy futures," *Preventing Chronic Disease* 4(3), July 3, 2007. See also, Washington State Department of Health, Tobacco Prevention and Control Program, Progress Report, March 2007. See also, Behavioral Risk Factor Surveillance System for adult smoking rates.
- ¹⁹ Youth Tobacco Surveillance in New York State, 2006. NY State Department of Health. http://www.nyhealth.gov/prevention/tobacco_control/youth_tobacco_survey.htm
- ²⁰ CDC, *Behavioral Risk Factor Surveillance System* (BRFSS), <http://apps.nccd.cdc.gov/brfss/list.asp?cat=TU&yr=2006&qkey=4396&state=All>.
- ²¹ *Maine 2007 Youth Risk Behavior Survey*, Maine Department of Human Services, 2008.
- ²² Indiana Youth Tobacco Survey, 2000 and 2006, http://www.in.gov/itpc/files/youth_fact_sheet_Jan08.pdf
- ²³ McAlister, AL, et al., "Settlement-Funded Tobacco Control in Texas: 2000-2004 Pilot Project Effects on Cigarette Smoking," *Public Health Reports*, May-June, 2006.
- ²⁴ Abt Associates Inc, *Independent Evaluation of the Massachusetts Tobacco Control Program, Seventh Annual Report, January 1994 to June 2000*.
- ²⁵ *Massachusetts Youth Risk Behavior Survey 2001; National Youth Risk Behavior Survey*.
- ²⁶ Abt Associates Inc, *Seventh Annual Report - January 1994 to June 2000*.
- ²⁷ National Health Interview Survey, 1993 and 2000.
- ²⁸ Abt Associates Inc, *Seventh Annual Report - January 1994 to June 2000*.
- ²⁹ Sbarra, C, Massachusetts Association of Health Boards, Abstract, March 2004. <http://www.mahb.org/tobacco/sales%20to%20minors%20study%20abstract.pdf>
- ³⁰ Data from Orzechowski & Walker, *Tax Burden on Tobacco 2006* [an industry-funded report]. Per capita cigarette consumption is measured as per capita cigarette pack sales.